



Not an actual patient.



ENHERTU[®]

fam-trastuzumab deruxtecan-nxki
20 mg/mL INJECTION FOR INTRAVENOUS USE

**FOR DISCUSSION WITH YOUR
HEALTHCARE PROVIDER:**

UNDERSTANDING AND MANAGING SIDE EFFECTS OF ENHERTU

Visit [ENHERTU.com/gastric](https://www.enhertu.com/gastric)
for more information

What is ENHERTU?

ENHERTU is a prescription medicine used in adults to treat human epidermal growth factor receptor 2 (HER2)-positive stomach cancer called gastric or gastroesophageal junction (GEJ) adenocarcinoma that has spread to areas near your stomach (locally advanced) or that has spread to other parts of your body (metastatic), and who have received a prior trastuzumab-based regimen.

It is not known if ENHERTU is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ENHERTU?

ENHERTU can cause serious side effects, including: Lung problems that may be severe, life-threatening or that may lead to death. If you develop lung problems your healthcare provider may treat you with corticosteroid medicines. Tell your healthcare provider right away if you get any of the following signs and symptoms:

- Cough
- Trouble breathing or shortness of breath
- Fever
- Other new or worsening breathing symptoms (e.g., chest tightness, wheezing)

Please see additional Important Safety Information on pages 15-19 of this brochure, and [click here for full ENHERTU Prescribing Information](#), including Boxed WARNINGS, and [click here for Medication Guide](#).

What is ENHERTU?

ENHERTU is a prescription medicine used in adults to treat human epidermal growth factor receptor 2 (HER2)-positive stomach cancer called gastric or gastroesophageal junction (GEJ) adenocarcinoma that has spread to areas near your stomach (locally advanced) or that has spread to other parts of your body (metastatic), and who have received a prior trastuzumab-based regimen.

It is not known if ENHERTU is safe and effective in children.

Find more information about ENHERTU
by visiting [ENHERTU.com/gastric](https://www.enhertu.com/gastric)

Important Safety Information

What is the most important information I should know about ENHERTU?

ENHERTU can cause serious side effects, including:

Lung problems that may be severe, life-threatening or that may lead to death. If you develop lung problems your healthcare provider may treat you with corticosteroid medicines. Tell your healthcare provider right away if you get any of the following signs and symptoms:

- Cough
- Trouble breathing or shortness of breath
- Fever
- Other new or worsening breathing symptoms (e.g., chest tightness, wheezing)

Please see additional Important Safety Information on pages 15-19 of this brochure, and [click here for full ENHERTU Prescribing Information](#), including Boxed WARNINGS, and [click here for Medication Guide](#).

How will I receive ENHERTU?

ENHERTU is given once every 3 weeks by intravenous (IV) infusion by your healthcare provider. Your healthcare provider will decide how many treatments you need.



The first infusion will take about 90 minutes so the doctor can see if there are any side effects or problems



Future infusions should take about 30 minutes, if your first infusion was well tolerated

- Your healthcare provider will give you medicines before your infusion to help prevent nausea and vomiting
- Your healthcare provider may slow down or temporarily stop your infusion of ENHERTU if you have an infusion-related reaction
- Your healthcare provider may permanently stop ENHERTU if you have severe infusion reactions
- If you miss a planned dose of ENHERTU, call your healthcare provider right away to schedule an appointment. Do not wait until the next planned treatment cycle

Important Safety Information (cont'd)

Low white blood cell counts (neutropenia). Low white blood cell counts are common with ENHERTU and can sometimes be severe. Your healthcare provider will check your white blood cell counts before starting ENHERTU and before starting each dose. Tell your healthcare provider right away if you develop any signs or symptoms of an infection or have fever or chills during treatment with ENHERTU.

 **ENHERTU**[®]
fam-trastuzumab deruxtecan-nxki
20 mg/mL INJECTION FOR INTRAVENOUS USE

What is the most important information you should know about ENHERTU?

ENHERTU can cause serious side effects.

Some serious or life-threatening side effects may affect your lungs, heart, or white blood cell count, affecting your ability to fight infection.

Pay special attention to new or worsening symptoms as they may be related to:




Lung problems that may be severe, life-threatening, or that may lead to death

- Tell your healthcare provider right away if you get any of the following signs and symptoms:
 - Cough
 - Trouble breathing or shortness of breath
 - Fever
 - Other new or worsening breathing symptoms (e.g., chest tightness, wheezing)

If you develop lung problems your healthcare provider may treat you with corticosteroid medicines.

Please see additional Important Safety Information on pages 15-19 of this brochure, and [click here for full ENHERTU Prescribing Information](#), including Boxed WARNINGS, and [click here for Medication Guide](#).



Not actual patients.



Low white blood cell counts (neutropenia)

- Low white blood cell counts are common with ENHERTU and can sometimes be severe
- Your healthcare provider will check your white blood cell counts before starting ENHERTU and before starting each dose
- Tell your healthcare provider right away if you develop any signs or symptoms of an infection or have fever or chills during treatment with ENHERTU

Continued on next page

Call or see your healthcare provider right away if you develop any symptoms of these side effects or if these symptoms get worse

 **ENHERTU**[®]
fam-trastuzumab deruxtecan-nxki
20 mg/mL INJECTION FOR INTRAVENOUS USE

What is the most important information you should know about ENHERTU? (cont'd)



Heart problems that may affect your heart's ability to pump blood

Your healthcare provider will check your heart function before starting treatment with ENHERTU.

Tell your healthcare provider right away if you get any of the following signs and symptoms:

- New or worsening shortness of breath
- Coughing
- Feeling tired
- Swelling of your ankles or legs
- Irregular heartbeat
- Sudden weight gain
- Dizziness or feeling light-headed
- Loss of consciousness

Getting medical treatment right away may help keep these side effects from becoming more serious



Harm to your unborn baby

- Tell your healthcare provider right away if you become pregnant or think you might be pregnant during treatment with ENHERTU
- If you are able to become pregnant, your healthcare provider should do a pregnancy test before you start treatment with ENHERTU
- **Females** who are able to become pregnant should use effective birth control (contraception) during treatment with ENHERTU and for 7 months after the last dose
- **Males** who have female partners that are able to become pregnant should use effective birth control (contraception) during treatment with ENHERTU and for 4 months after the last dose

Your healthcare provider will check you for side effects during your treatment with ENHERTU. Tell your healthcare provider about any side effects you may experience. Your healthcare provider may reduce your dose, delay treatment or completely stop treatment with ENHERTU if you have severe side effects

What are the possible side effects of ENHERTU?

ENHERTU can cause serious side effects. Please see pages 4-7 and 15-19 for more information.

The most common side effects of ENHERTU, when used to treat stomach cancer, include:

- Low red blood cell counts
- Low white blood cell counts
- Low platelet counts
- Nausea
- Decreased appetite
- Increased liver function tests
- Feeling tired
- Diarrhea
- Low levels of blood potassium
- Vomiting
- Constipation
- Fever
- Hair loss

ENHERTU may cause fertility problems in males, which may affect the ability to father children. Talk to your healthcare provider if you have concerns about fertility.

These are not all of the possible side effects of ENHERTU. Tell your healthcare provider if you experience any side effects, including those not listed in this guide.

Call your healthcare provider for medical advice about side effects.

You are encouraged to report negative side effects of ENHERTU by calling 1-877-437-7763.

If you prefer to report these to the FDA, visit [FDA.gov/medwatch](https://www.fda.gov/medwatch) or call 1-800-FDA-1088 (1-800-332-1088)

Please see additional Important Safety Information on pages 15-19 of this brochure, and [click here for full ENHERTU Prescribing Information](#), including Boxed WARNINGS, and [click here for Medication Guide](#).

Let your healthcare provider know right away if you are experiencing any side effects

Your healthcare provider will check you for these side effects during your treatment with ENHERTU. Your healthcare provider may reduce your dose, delay treatment or completely stop treatment with ENHERTU if you have severe side effects.

Talk to your healthcare provider before taking any medications (including prescription or over-the-counter medicines, vitamins, and herbal supplements) or making changes to your diet or daily activities

 **ENHERTU[®]**
fam-trastuzumab deruxtecan-nxki
20 mg/mL INJECTION FOR INTRAVENOUS USE

Useful tips to help you manage some of these side effects



Nausea and vomiting

- Try to eat 5 or 6 small meals and snacks each day instead of 3 large meals
- Eat bland, easy-to-digest food and drinks
- Avoid food and drinks that are too hot or too cold
- Avoid strong smells if possible
- If you feel like vomiting, take deep, slow breaths and, if possible, get fresh air

Important Safety Information (cont'd)

Heart problems that may affect your heart's ability to pump blood. Your healthcare provider will check your heart function before starting treatment with ENHERTU. Tell your healthcare provider right away if you get any of the following signs and symptoms:

- New or worsening shortness of breath
- Coughing
- Feeling tired
- Swelling of your ankles or legs
- Irregular heartbeat
- Sudden weight gain
- Dizziness or feeling light-headed
- Loss of consciousness

Please see additional Important Safety Information on pages 15-19 of this brochure, and [click here for full ENHERTU Prescribing Information](#), including **Boxed WARNINGS**, and [click here for Medication Guide](#).



Diarrhea

- Try to eat 5 or 6 small meals and snacks each day instead of 3 large meals
- Eat foods low in fiber
- Drink clear liquids to replace lost fluids. Check with your healthcare provider for how much fluid you should drink and what types of fluid might be best for you

Talk to your healthcare provider about these tips and before taking any medicine for nausea or diarrhea

Important Safety Information (cont'd)

Your healthcare provider will check you for these side effects during your treatment with ENHERTU. Your healthcare provider may reduce your dose, delay treatment or completely stop treatment with ENHERTU if you have severe side effects.

 **ENHERTU**[®]
fam-trastuzumab deruxtecan-nxki
20 mg/mL INJECTION FOR INTRAVENOUS USE

Useful tips to help you manage some of these side effects (cont'd)



Feeling tired (fatigue)

- Talk to your healthcare provider about what activities may be right for you
- Meditation may help with relaxation and decrease stress
- Plan rest time and try not to do too much
- Try to get regular exercise during times when your energy level is high (according to your healthcare provider's approval)
- Let other people help you when you are feeling tired. They can assist with household chores, shopping, cooking, or driving to and from doctor's visits

Talk to your healthcare provider to help determine the cause of why you may be feeling tired

Important Safety Information (cont'd)

Harm to your unborn baby. Tell your healthcare provider right away if you become pregnant or think you might be pregnant during treatment with ENHERTU.

- If you are able to become pregnant, your healthcare provider should do a pregnancy test before you start treatment with ENHERTU.

Please see additional Important Safety Information on pages 15-19 of this brochure, and [click here for full ENHERTU Prescribing Information](#), including Boxed WARNINGS, and [click here for Medication Guide](#).



Dealing with hair thinning or hair loss (alopecia)

Before hair loss or thinning:

- Talk to your healthcare provider about the possibility of hair loss
- Consider cutting your hair short to feel more in control of hair loss
- Be gentle when you wash your hair. Use a mild shampoo, such as baby shampoo
- Consider asking your healthcare provider about cold caps or scalp cooling systems. These may help reduce hair loss by narrowing the blood vessels beneath your scalp
- Avoid items such as electric hair dryers, hairspray, hair dyes, products to perm or relax hair, and hair bands and clips that can hurt your scalp
- If you plan on buying a wig, you should do so while you still have hair

If hair loss or thinning occurs:

- Protect your scalp
- Cover your head to help stay warm. You may feel colder once you lose your hair
- Sleep on a satin pillowcase. Satin creates less friction and may be more comfortable

If you are concerned about hair loss, you might want to talk with a member of your healthcare team, a friend or family member, or another person like you undergoing treatment

 **ENHERTU**[®]
fam-trastuzumab deruxtecan-nxki
20 mg/mL INJECTION FOR INTRAVENOUS USE

What should I tell my healthcare provider before starting ENHERTU?

Tell your healthcare provider about all of your medical conditions, including if you:

- Have lung or breathing problems
 - Have signs or symptoms of an infection
 - Have or have had any heart problems
 - Have kidney problems. Your healthcare provider may follow you more closely. In clinical trials, more serious lung problems were seen in patients with certain kidney problems
 - Have liver problems. Your healthcare provider may follow you more closely
 - Are breastfeeding or plan to breastfeed
- It is not known if ENHERTU passes into your breast milk. Do not breastfeed during treatment with ENHERTU and for 7 months after the last dose

Always let your healthcare provider know about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements

Important Safety Information

What is the most important information I should know about ENHERTU?

ENHERTU can cause serious side effects, including:

Lung problems that may be severe, life-threatening or that may lead to death.

If you develop lung problems your healthcare provider may treat you with corticosteroid medicines. Tell your healthcare provider right away if you get any of the following signs and symptoms:

- Cough
- Trouble breathing or shortness of breath
- Fever
- Other new or worsening breathing symptoms (e.g., chest tightness, wheezing)

Low white blood cell counts

(neutropenia). Low white blood cell counts are common with ENHERTU and can sometimes be severe. Your healthcare provider will check your white blood cell counts before starting ENHERTU and before starting each dose. Tell your healthcare provider right away if you develop any signs or symptoms of an infection or have fever or chills during treatment with ENHERTU.

Continued on next page

Important Safety Information (cont'd)

Heart problems that may affect your heart's ability to pump blood. Your healthcare provider will check your heart function before starting treatment with ENHERTU. Tell your healthcare provider right away if you get any of the following signs and symptoms:

- New or worsening shortness of breath
- Coughing
- Feeling tired
- Swelling of your ankles or legs
- Irregular heartbeat
- Sudden weight gain
- Dizziness or feeling light-headed
- Loss of consciousness

Your healthcare provider will check you for these side effects during your treatment with ENHERTU. Your healthcare provider may reduce your dose, delay treatment or completely stop treatment with ENHERTU if you have severe side effects.

Harm to your unborn baby. Tell your healthcare provider right away if you become pregnant or think you might be pregnant during treatment with ENHERTU.

- If you are able to become pregnant, your healthcare provider should do a pregnancy test before you start treatment with ENHERTU.

- **Females** who are able to become pregnant should use effective birth control (contraception) during treatment with ENHERTU and for 7 months after the last dose.
- **Males** who have female partners that are able to become pregnant should use effective birth control (contraception) during treatment with ENHERTU and for 4 months after the last dose.

Before you receive ENHERTU, tell your healthcare provider about all of your medical conditions, including if you:

- Have lung or breathing problems.
- Have signs or symptoms of an infection.
- Have or have had any heart problems.
- Are breastfeeding or plan to breastfeed. It is not known if ENHERTU passes into your breast milk. Do not breastfeed during treatment with ENHERTU and for 7 months after the last dose.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will I receive ENHERTU?

- You will receive ENHERTU into your vein through an intravenous (IV) line by your healthcare provider.

Continued on next page

Important Safety Information (cont'd)

- ENHERTU is given 1 time every three weeks (21-day treatment cycle).
- Your healthcare provider will decide how many treatments you need.
- Your healthcare provider will give you medicines before your infusion to help prevent nausea and vomiting.
- Your healthcare provider may slow down or temporarily stop your infusion of ENHERTU if you have an infusion-related reaction, or permanently stop ENHERTU if you have severe infusion reactions.
- If you miss a planned dose of ENHERTU, call your healthcare provider right away to schedule an appointment. Do not wait until the next planned treatment cycle.

What are the possible side effects of ENHERTU?

ENHERTU can cause serious side effects. See “What is the most important information I should know about ENHERTU?”

The most common side effects of ENHERTU, when used in people with HER2-positive gastric or GEJ adenocarcinoma, include:

- Low red blood cell counts
- Low white blood cell counts
- Low platelet counts
- Nausea
- Decreased appetite
- Increased liver function tests
- Feeling tired
- Diarrhea
- Low levels of blood potassium
- Vomiting
- Constipation
- Fever
- Hair loss

Please see additional Important Safety Information on pages 15-19 of this brochure, and [click here for full ENHERTU Prescribing Information](#), including **Boxed WARNINGS**, and [click here for Medication Guide](#).

ENHERTU may cause fertility problems in males, which may affect the ability to father children. Talk to your healthcare provider if you have concerns about fertility.

These are not all of the possible side effects of ENHERTU. Call your doctor for medical advice about side effects. You may report side effects to Daiichi Sankyo at 1-877-437-7763 or to FDA at 1-800-FDA-1088.

What is ENHERTU?

ENHERTU is a prescription medicine used in adults to treat human epidermal growth factor receptor 2 (HER2)-positive stomach cancer called gastric or gastroesophageal junction (GEJ) adenocarcinoma that has spread to areas near your stomach (locally advanced) or that has spread to other parts of your body (metastatic), and who have received a prior trastuzumab-based regimen.

It is not known if ENHERTU is safe and effective in children.

Please see full Prescribing Information, including Boxed WARNINGS, and Medication Guide.

 **ENHERTU**[®]
fam-trastuzumab deruxtecan-nxki
20 mg/mL INJECTION FOR INTRAVENOUS USE



Not actual patients.

Find out more about ENHERTU
**Talk to your healthcare
provider and visit**
[ENHERTU.com/learnmore](https://www.enhertu.com/learnmore)
or scan the QR code



What is ENHERTU?

ENHERTU is a prescription medicine used in adults to treat human epidermal growth factor receptor 2 (HER2)-positive stomach cancer called gastric or gastroesophageal junction (GEJ) adenocarcinoma that has spread to areas near your stomach (locally advanced) or that has spread to other parts of your body (metastatic), and who have received a prior trastuzumab-based regimen.

It is not known if ENHERTU is safe and effective in children.

References: **1.** ENHERTU [prescribing information]. Daiichi Sankyo Inc., Basking Ridge, NJ and AstraZeneca Pharmaceuticals LP, Wilmington, DE, 2022. **2.** National Cancer Institute. Chemotherapy and you. Accessed June 8, 2022. <https://www.cancer.gov/publications/patient-education/chemotherapy-and-you.pdf>. **3.** American Cancer Society. Cancer.org. Accessed August 9, 2022. <https://www.cancer.org/treatment/treatments-and-side-effects/physical-side-effects/hair-loss/cold-caps.html>.

Please see additional Important Safety Information on pages 15-19 of this brochure, and [click here for full ENHERTU Prescribing Information](#), including **Boxed WARNINGS, and [click here for Medication Guide](#).**

 **ENHERTU**[®]
fam-trastuzumab deruxtecan-nxki
20 mg/mL INJECTION FOR INTRAVENOUS USE

 Daiichi-Sankyo | AstraZeneca 

ENHERTU[®] is a registered trademark of Daiichi Sankyo Company, Limited.
© 2022 Daiichi Sankyo, Inc. and AstraZeneca. PP-US-ENG-0683 10/22